

JEMDSA

Journal of Endocrinology, Metabolism and Diabetes of South Africa

JEMDSA 2025 Volume 30 Number 2

Please answer the following questionnaire online.

SEMDSA and other Society members:

1. Visit our new CPD portal at <https://www.mpconsulting.co.za/>
2. Create a new account and set your CPD requirements.
3. Not a new user? You may have been registered with the following details: username: e-mail address and password: MP+0+6digits (e.g. MP0123456). All records from the previous system have now been moved to the new platform.
4. For CPD assistance please phone 086 1111 335

1. What is the effect of a 1 mmol/L reduction in LDL on major coronary events in people with type 2 diabetes?

- a. 4% reduction
- b. 14% reduction
- c. 24% reduction
- d. 34% reduction

2. What is the target LDL value for primary prevention in people with type 2 diabetes?

- a. < 1.2 mmol/L
- b. < 1.8 mmol/L
- c. < 2.6 mmol/L
- d. < 3.0 mmol/L

3. What is the prevalence of silent myocardial ischaemia in people with diabetes?

- a. 0.8%
- b. 8.5%
- c. 18.5%
- d. 28.5%

4. Hypogonadism and other causes of male sexual dysfunction are uncommon in men with diabetes

- a. True
- b. False

5. What is the most common cause of primary hyperparathyroidism?

- a. single adenoma
- b. bilateral adenomas
- c. hyperplasia
- d. carcinoma