

# The evolution of obesity pharmacotherapy from sympathomimetics to incretin-based therapies

KN Ncube,<sup>1</sup> NL Moloi,<sup>1</sup> TD Malange<sup>2</sup>

<sup>1</sup> Department of Pharmacology, Faculty of Health Sciences, University of Pretoria, South Africa

<sup>2</sup> Department of Clinical Pharmacology, Faculty of Medicine and Health Sciences, Stellenbosch University, South Africa

Corresponding author, email: keith.ncube@up.ac.za

## Abstract

Obesity is a chronic metabolic disorder that has reached epidemic proportions globally and in South Africa, contributing to the increasing burden of cardiometabolic diseases. Although lifestyle modifications remain a fundamental approach, long-term weight loss is often limited, necessitating the use of pharmacotherapy. Historically, centrally acting sympathomimetics, such as phentermine, have been predominant in South African treatments, albeit with restrictions concerning their safety and duration. This review examines the evolution of obesity pharmacotherapy from traditional agents, including phentermine and orlistat, to contemporary incretin-based therapies. Particular emphasis is placed on glucagon-like peptide-1 receptor agonists and dual incretin agonists, such as semaglutide and tirzepatide, which have demonstrated unprecedented efficacy in clinical trials involving patients with obesity. Emerging multi-hormonal and non-injectable agents are also discussed. This article underscores the transition from short-term appetite suppression to sustained pharmacological management of obesity and its comorbidities.

**Keywords:** incretin-based therapies, obesity, pharmacotherapy, weight management

© Authors

<https://doi.org/10.36303/SAPJ.4108>

## Introduction

Obesity has reached epidemic proportions worldwide. There has been a startling rise in obesity rates, with prevalence having tripled since 1975 and now affecting over 13% of the global adult population.<sup>1</sup> This rapid increase, together with the rising burden of obesity-related comorbidities, has intensified the need for more effective therapeutic strategies.<sup>1</sup> Obesity is commonly defined as an accumulation of abnormal or excessive adipose tissue associated with an increase in Body Mass Index (BMI).<sup>2</sup> The BMI is a calculation that utilises a person's weight in kilograms divided by their height in meters. The World Health Organization (WHO) proposes that obesity and morbid obesity are proportional to a BMI of  $\geq 30$  kg/m<sup>2</sup> and  $\geq 40$  kg/m<sup>2</sup>, respectively.<sup>3</sup> Although this calculation is ubiquitously used, it is not without pitfalls. The BMI calculation does not account for body composition, as it does not consider other tissues such as muscle, and can thus under- or overestimate adiposity.<sup>4</sup> Despite contention of the usefulness of the index, it is currently the surrogate to determine risks associated with DM and other metabolic illnesses. It can be argued that abdominal adiposity would be a more effective indicator as it is linked with insulin resistance and other metabolic disorders.<sup>5</sup> Some experts propose that obesity should be viewed as a disease, but no consensus has been reached on this matter.<sup>4</sup>

The rising prevalence of obesity reflects the complex interplay of environmental, behavioural, and socioeconomic factors. As a result of urbanisation and evolving socioeconomic dynamics, processed foods have become more popular, and these contain high levels of refined sugars, flavouring agents, and emulsifiers.<sup>6</sup> These additives have demonstrated addictive properties,<sup>6,7</sup>

inferring repetitive use. Multiple studies have indicated a strong association between these factors and obesity. Numerous observational and experimental studies have demonstrated a strong association between these dietary patterns and excess weight gain.<sup>6</sup> Clinicians have been managing obesity and its sequelae for years. However, despite these efforts, it is now the fifth leading cause of mortality globally.<sup>3</sup> Care and prevention have mainly revolved around lifestyle modifications, but these are less effective in isolation.<sup>8</sup> Additionally, severely obese patients have mentioned having difficulty implementing changes owing to their environment and schedules.<sup>9</sup>

## Pathophysiology of obesity

The aetiology of obesity is multifactorial, involving genetic and environmental factors that ultimately lead to sustained dysregulation of energy homeostasis.<sup>10</sup> While single-gene defects or syndromic forms contribute to rare cases of paediatric obesity, cases arise from a complex interplay between inherited susceptibility and environmental influences.<sup>10</sup> Central regulation of appetite is disrupted within the arcuate nucleus of the hypothalamus, where an imbalance between orexigenic agouti-related peptide and neuropeptide Y neurons and anorexigenic pro-opiomelanocortin and cocaine and amphetamine-regulated transcript neurons impairs satiety signalling.<sup>11</sup> Normally, these hypothalamic satiety signalling pathways are reinforced by gut-derived incretin hormones such as glucagon-like peptide 1 (GLP-1) and the glucose-dependent insulinotropic polypeptide (GIP).<sup>12-13</sup> In obesity, responsiveness to these signals is attenuated, contributing to persistent appetite drive.<sup>12</sup> Concurrently, sustained positive energy balance promotes adipocyte hypertrophy

and hyperplasia, leading to the expansion of adipose tissue.<sup>14</sup> This expansion disrupts adipokine secretion, leading to high leptin levels and the development of central leptin resistance.<sup>14</sup> Additionally, the release of pro-inflammatory cytokines<sup>15</sup> and activation of the endocannabinoid system<sup>16</sup> drive chronic low-grade inflammation and insulin resistance. Together, these neurohormonal and metabolic shifts render obesity a chronic disease in which the body's internal systems actively maintain an elevated baseline weight. Pharmacological approaches to obesity often overlap with those used in type 2 diabetes mellitus.<sup>17</sup>

**Pharmacological therapies**

Lifestyle measures, including dietary modification, physical activity, and behavioural interventions, achieve modest initial weight loss and improve metabolic health.<sup>18</sup> However, long-term maintenance is often hindered by adaptive thermogenesis, increased appetite, and the practical difficulty of sustaining habits in daily life.<sup>19</sup> Consequently, pharmacotherapy plays a vital role as an adjunct to lifestyle modifications in the treatment of obesity, especially for patients who do not achieve sufficient weight loss through lifestyle interventions alone.<sup>20</sup> A variety of centrally acting, peripherally acting, and incretin-based agents are currently available or in development, each with distinct mechanisms of action and efficacy profiles, as shown in Table 1.

**Centrally acting sympathomimetic appetite suppressants**

Centrally acting drugs, such as phentermine (Duromine®), have long been used as adjuncts in the management of obesity. These agents work by increasing the concentrations of norepinephrine, dopamine, and serotonin within the hypothalamic nuclei, consequently enhancing satiety and suppressing appetite. Phentermine was approved by the United States Food and Drug Administration in 1959 for short-term weight management, with treatment duration limited to 12 weeks, though it remains used

off-label long-term. In South Africa, Duromine® is registered for short-term obesity management in doses of 15 or 30 mg daily.<sup>20</sup> Phentermine monotherapy is associated with modest short-term weight loss, with controlled trials reporting reductions of approximately 3–5% of baseline body weight. A quantitative analysis of clinical trials of phentermine monotherapy revealed a mean weight loss of 3.6 kg over a mean of 13 weeks.<sup>21</sup> Common side-effects of phentermine include tachycardia, restlessness, insomnia, dry mouth and constipation.<sup>30</sup> Despite weight loss typically leading to reduced blood pressure, phentermine can increase blood pressure and is thus not recommended for people with uncontrolled hypertension.<sup>30</sup> To enhance efficacy, phentermine is commonly combined with topiramate, which modulates γ-aminobutyric acid (GABA) receptors, synergistically increasing satiety.<sup>22</sup> The landmark CONQUER trial demonstrated that this dual approach achieves a superior 10.2% mean weight loss over 56 weeks, nearly doubling the efficacy of monotherapy, while allowing for lower, better-tolerated dosing for long-term obesity management.<sup>22</sup>

**Centrally acting combination therapies**

Combination therapies that target multiple central pathways involved in appetite regulation include the naltrexone-bupropion combination, marketed as Contrave. Approved in South Africa in 2020, this fixed-dose combination contains 16 mg of naltrexone and 180 mg of bupropion per tablet, administered twice daily.<sup>20</sup> Naltrexone is an opioid receptor antagonist that inhibits β-endorphin-mediated autoinhibitory feedback of pro-opiomelanocortin neurons, thereby sustaining satiety signals. Conversely, bupropion functions as a dopamine and norepinephrine reuptake inhibitor that stimulates these neurons and further enhances satiety. In the COR-I trial, treatment over 56 weeks resulted in a mean weight loss of 6.1% compared to 1.3% in the placebo group, with 48% of participants achieving at least

**Table 1:** Pharmacological agents used for the management of obesity

Drug/ Combination	Drug class	Primary mechanism of action	Typical dosing	Reference(s)
Phentermine (Duromine®)	Centrally acting sympathomimetic	Increases hypothalamic norepinephrine, dopamine, and serotonin to suppress appetite	15–30 mg once daily	21
Phentermine + Topiramate	Centrally acting combination	Sympathomimetic appetite suppression plus GABA modulation	Titrated combination dosing	22
Naltrexone + Bupropion (Contrave®)	Centrally acting combination	POMC activation and inhibition of β-endorphin feedback	16 mg/180 mg twice daily	23
Orlistat (Xenical®)	Peripheral lipase inhibitor	Inhibits gastric and pancreatic lipases, reducing fat absorption	120 mg three times daily	24
Liraglutide (Saxenda®)	GLP-1 receptor agonist	Enhances satiety; delays gastric emptying; improves glycaemic control	3.0 mg once daily (SC)	25
Semaglutide (Wegovy®)	GLP-1 receptor agonist	Central appetite suppression and peripheral metabolic effects	2.4 mg once weekly (SC)	26
Tirzepatide (Mounjaro®)	Dual GLP-1/GIP receptor agonist	Multi-hormonal appetite and metabolic regulation	Weekly SC dosing (titrated)	27
Emerging agents (e.g. retatrutide, orforglipron, cagrilintide)	Multi-agonists/novel incretins	Target multiple metabolic and appetite pathways	Under investigation	28,29

Abbreviations: GABA: gamma-aminobutyric acid; GI: gastrointestinal; GIP: glucose-dependent insulinotropic polypeptide; GLP-1: glucagon-like peptide-1; SC: subcutaneous

5.0% weight loss.<sup>23</sup> Common adverse effects include nausea, constipation, and headache. Owing to the bupropion component, this combination is contraindicated in patients with seizure disorders or uncontrolled hypertension, and it should not be co-administered with opioids.<sup>20</sup>

### Peripheral gastrointestinal lipase inhibitors

Peripheral gastrointestinal lipase inhibitors decrease fat absorption in the gastrointestinal tract, with orlistat (Xenical®) representing the main clinically used agent. Registered in South Africa since 1997, orlistat is a potent inhibitor of gastric and pancreatic lipases and functions by binding covalently to the active serine residues of these enzymes in the gastric and intestinal lumen.<sup>20</sup> Unlike central-acting drugs, orlistat acts peripherally to inhibit the hydrolysis and absorption of dietary triglycerides, leading to a subsequent faecal excretion of approximately 30% of ingested fat.<sup>31</sup> Clinical evidence from a recent systematic review of 22 randomised controlled trials indicates that orlistat, taken at a dose of 120 mg three times daily, results in a mean weight loss of approximately 2.4 kg greater than placebo.<sup>32</sup> Beyond its role in weight reduction, the landmark XENDOS trial has demonstrated that orlistat, when administered as an adjunct to lifestyle intervention, significantly reduces the cumulative incidence of type 2 diabetes by 37.3% in patients with impaired glucose tolerance.<sup>24</sup> However, the clinical utility of orlistat is often limited by gastrointestinal side-effects due to fat malabsorption. Such side-effects include steatorrhoea, oily spotting, faecal urgency, diarrhoea, abdominal pain, and fatty/oily stools.<sup>33</sup> Additionally, orlistat can impair the absorption of fat-soluble vitamins A, D, E, and K, occasionally leading to reduced plasma concentrations; thus, supplementation with a multivitamin is recommended during therapy to prevent deficiencies.<sup>33</sup>

### Incretin-based therapies and metabolic appetite regulation

Incretin-based therapies have expanded the pharmacological options available for obesity management by mimicking endogenous hormones involved in the regulation of satiety and glucose homeostasis.<sup>34</sup> Liraglutide and semaglutide are the most commonly used agents within this class, and they are glucagon-like peptide-1 receptor agonists (GLP-1 RAs) that target the hypothalamic pro-opiomelanocortin and cocaine- and amphetamine-regulated transcript neurons, thereby enhancing satiety and reducing hunger.<sup>35</sup> Peripherally, these agents slow gastric emptying and modulate glucose metabolism by stimulating glucose-dependent insulin secretion and suppressing glucagon release.<sup>35</sup> The SCALE trial demonstrated that liraglutide administered at a dose of 3 mg daily over 56 weeks results in a mean weight loss of approximately 8%.<sup>36</sup> Moreover, the more potent, once-weekly 2.4 mg semaglutide led to a mean weight loss of 14.9% in the STEP 1 trial.<sup>26</sup> Direct comparative data from the STEP 8 trial further demonstrated the superior efficacy of semaglutide, with a mean weight loss of 15.8% compared with 6.4% for liraglutide.<sup>37</sup> Despite their efficacy, GLP-1 RAs are commonly associated with dose-dependent gastrointestinal

side-effects, such as nausea and vomiting, which typically require gradual dose escalation to improve tolerability.<sup>35</sup>

### Emerging therapies and combination regimens

Obesity pharmacotherapy is rapidly developing, with a shift towards the use of multi-hormonal co-agonists that target multiple pathways to achieve weight loss comparable to that achieved with bariatric surgery. Tirzepatide, a dual GLP-1 and GIP receptor agonist, demonstrated a mean weight loss of up to 22.5% at the highest dose in the SURMOUNT-1 trial,<sup>27</sup> leading to its recent approval for obesity management in South Africa.<sup>20</sup> Beyond dual agonism, triple agonists such as retatrutide which targets the GLP-1, GIP, and glucagon receptors, have shown unprecedented efficacy. In a Phase 2 trial, participants achieved a mean weight loss of 24.2% at 48 weeks,<sup>38</sup> and subsequent Phase 3 TRIUMPH-4 data reported to achieve weight loss of 28.7% over 68 weeks.<sup>39</sup>

There is also a renewed interest in treating obesity by modulating the endocannabinoid system through the use of peripherally restricted cannabinoid-1 (CB1) receptor blockers such as monlunabant and nimacimab.<sup>40-41</sup> Unlike the first-generation CB-1 receptor antagonist rimonabant,<sup>42</sup> which was withdrawn due to psychiatric risks, these newer agents are designed to target metabolic tissues while avoiding the central nervous system, thus minimising neuropsychiatric side-effects.<sup>43</sup> Additionally, the development of oral incretins such as orforglipron<sup>28</sup> and amylin-based combinations such as cagrilintide<sup>29</sup> indicates that there is an emerging focus on highly potent, scalable, and non-injectable options for long-term obesity management.

### Conclusion and future directions

Obesity is a chronic metabolic disorder that requires a comprehensive therapeutic approach. In South Africa, pharmacological management has relied on the short-term use of centrally acting sympathomimetics, such as phentermine, which offers modest efficacy but is limited by safety concerns. The advent of more effective incretin-based therapies marks a significant advance in obesity pharmacotherapy, providing durable weight reduction and metabolic improvements. These agents have redefined therapeutic expectations by enabling sustained control of appetite and energy balance rather than temporary weight loss. As the therapeutic landscape evolves to include dual and triple hormonal agonists, clinical practice must align with emerging evidence and guidelines. The integration of pharmacotherapy with lifestyle interventions remains crucial for effective obesity management and reducing obesity-related morbidity.

### ORCID

KN Ncube  <https://orcid.org/0000-0003-4693-9868>

NL Moloi  <https://orcid.org/0009-0003-1083-3006>

TD Malange  <https://orcid.org/0000-0002-4963-1386>

### References

1. Ahmed SK, Mohammed RA. Obesity: Prevalence, causes, consequences, management, preventive strategies and future research directions. *Metabolism Open*. 2025;27:100375.

- <https://doi.org/10.1016/j.metop.2025.100375>.
- May W, Goedecke JH, Conradi-Smit M. The science of obesity. *S Afr Med J*. 2025; 115(10b):e3600. <https://doi.org/10.7196/SAMJ.2025.v115i9b.3600>.
  - World Health Organization. Obesity and overweight. World Health Organization (WHO); 2025 Available from: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>. Accessed 09 January 2026.
  - Rubino F, Cummings DE, Eckel RH, et al. Definition and diagnostic criteria of clinical obesity. *Lancet Diabetes & Endocrinology*. 2025;13(3):221-62. [https://doi.org/10.1016/S2213-8587\(24\)00316-4](https://doi.org/10.1016/S2213-8587(24)00316-4).
  - Rodriguez-Granillo GA, Carrascosa P. Abdominal obesity as a risk predictor. *JACC*. 2018;71(12):1398-9. <https://doi.org/10.1016/j.jacc.2017.12.065>.
  - Canhada SL, Luft VC, Giatti L, et al. Ultra-processed foods, incident overweight and obesity, and longitudinal changes in weight and waist circumference: The Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *Public Health Nutr*. 2020;23(6):1076-86. <https://doi.org/10.1017/S1368980019002854>.
  - Schiestl ET, Gearhardt AN, Wolfson J. The qualitative evaluation of food addiction across the lifespan. *Appetite*. 2024;194:107170. <https://doi.org/10.1016/j.appet.2023.107170>.
  - Fatoye F, Mbada C, Niyi-Odumosu F, et al. The clinical and economic burden of obesity in low- and middle-income countries: A systematic review. *Int J Obes*. 2025;49(12):2453-61. <https://doi.org/10.1038/s41366-025-01913-3>.
  - Toft BS, Hörberg U, Rasmussen B. The ups and downs of lifestyle modification: An existential journey among persons with severe obesity. *Scand J Caring Sci*. 2022;36(1):265-74. <https://doi.org/10.1111/scs.12985>.
  - Flores-Dorantes MT, Díaz-López YE, Gutiérrez-Aguilar R. Environment and gene association with obesity and its impact on neurodegenerative and neurodevelopmental diseases. *Front Neurosci*. 2020;14. <https://doi.org/10.3389/fnins.2020.00863>.
  - Vohra MS, Benchoula K, Serpell CJ, Hwa WE. AgRP/NPY and POMC neurons in the arcuate nucleus and their potential role in treatment of obesity. *Eur J Pharmacol*. 2022;915:174611. <https://doi.org/10.1016/j.ejphar.2021.174611>.
  - Ranganath LR, Beety JM, Morgan LM, et al. Attenuated GLP-1 secretion in obesity: Cause or consequence? *Gut*. 1996;38(6):916-9. <https://doi.org/10.1136/gut.38.6.916>.
  - Holst JJ, Rosenkilde MM. GIP as a therapeutic target in diabetes and obesity: Insight from incretin co-agonists. *Journal of Clinical Endocrinology & Metabolism*. 2020;105(8):e2710-e6. <https://doi.org/10.1210/clinem/dgaa327>.
  - Horwitz A, Birk R. Adipose tissue hyperplasia and hypertrophy in common and syndromic obesity—the case of BBS obesity. *Nutrients*. 2023;15(15):3445. <https://doi.org/10.3390/nu15153445>.
  - Turner L, Wanasinghe Anjalee I, Brunori P, Santosa S. Is adipose tissue inflammation the culprit of obesity-associated comorbidities? *Obes Rev*. 2025;26(11):e13956. <https://doi.org/10.1111/obr.13956>.
  - Gruden G, Barutta F, Kunos G, Pacher P. Role of the endocannabinoid system in diabetes and diabetic complications. *Br J Pharmacol*. 2016;173(7):1116-27. <https://doi.org/10.1111/bph.13226>.
  - Ncube K, Malange TD. Beyond metformin: The expanding landscape of type 2 diabetes treatment. *SA Pharmaceutical Journal*. 2025;92(1):30-4. <https://doi.org/10.36303/SAPJ.2299>.
  - Buchanan L, Calkins M, Kalayjian T, et al. Toward a metabolic health intervention, demonstrates robust 1-year weight loss and cost-savings through deprescription. *Frontiers in Nutrition*. 2025;12. <https://doi.org/10.3389/fnut.2025.1548609>.
  - Isola V, Hulmi JJ, Petäjä P, et al. Weight loss induces changes in adaptive thermogenesis in female and male physique athletes. *Applied Physiology, Nutrition, and Metabolism*. 2023;48(4):307-20. <https://doi.org/10.1139/apnm-2022-0372>.
  - Noeth M, Van Zyl F, Hellig J, Conradi-Smit M, May W. Pharmacotherapy for obesity management. *S Afr Med J*. 2025;115(9b):e3767-e. <https://doi.org/10.7196/SAMJ.2025.v115i9b.3767>.
  - Haddock CK, Poston WSC, Dill PL, Foreyt JP, Ericsson M. Pharmacotherapy for obesity: A quantitative analysis of four decades of published randomized clinical trials. *Int J Obes*. 2002;26(2):262-73. <https://doi.org/10.1038/sj.jco.0801889>.
  - Gadde KM, Allison DB, Ryan DH, et al. Effects of low-dose, controlled-release, phentermine plus topiramate combination on weight and associated comorbidities in overweight and obese adults (conquer): A randomised, placebo-controlled, phase 3 trial. *The Lancet*. 2011;377(9774):1341-52. [https://doi.org/10.1016/S0140-6736\(11\)60205-5](https://doi.org/10.1016/S0140-6736(11)60205-5).
  - Greenway FL, Fujioka K, Plodkowski RA, et al. Effect of naltrexone plus bupropion on weight loss in overweight and obese adults (COR-1): A multicentre, randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet*. 2010;376(9741):595-605. [https://doi.org/10.1016/S0140-6736\(10\)60888-4](https://doi.org/10.1016/S0140-6736(10)60888-4).
  - Torgerson JS, Hauptman J, Boldrin MN, Sjörström L. Xenical in the prevention of diabetes in obese subjects (xendos) study: A randomized study of orlistat as an adjunct to lifestyle changes for the prevention of type 2 diabetes in obese patients. *Diabetes Care*. 2004;27(1):155-61. <https://doi.org/10.2337/diacare.27.1.155>.
  - Mok J, Adeleke MO, Brown A, et al. Safety and efficacy of liraglutide, 3.0 mg, once daily vs placebo in patients with poor weight loss following metabolic surgery: The BARI-OPTIMISE randomized clinical trial. *JAMA Surgery*. 2023;158(10):1003-11. <https://doi.org/10.1001/jamasurg.2023.2930>.
  - Wilding JPH, Batterham RL, Calanna S, et al. Once-weekly semaglutide in adults with overweight or obesity. *N Engl J Med*. 2021;384(11):989-1002. <https://doi.org/10.1056/NEJMoa2032183>.
  - Jastreboff AM, Aronne LJ, Ahmad NN, et al. Tirzepatide once weekly for the treatment of obesity. *N Engl J Med*. 2022;387(3):205-16. <https://doi.org/10.1056/NEJMoa2206038>.
  - Wharton S, Aronne LJ, Stefanski A, et al. Orforglipron, an oral small-molecule glp-1 receptor agonist for obesity treatment. *N Engl J Med*. 2025;393(18):1796-806. <https://doi.org/10.1056/NEJMoa2511774>.
  - Carvas AO, Leuthardt A, Kulka P, et al. Cagrilintide lowers bodyweight through brain amylin receptors 1 and 3. *eBioMedicine*. 2025;118. <https://doi.org/10.1016/j.ebiom.2025.105836>.
  - Hendricks EJ, Greenway FL, Westman EC, Gupta AK. Blood pressure and heart rate effects, weight loss and maintenance during long-term phentermine pharmacotherapy for obesity. *Obesity*. 2011;19(12):2351-60. <https://doi.org/10.1038/oby.2011.94>.
  - Guercioli R. Mode of action of orlistat. *Int J Obes Relat Metab Disord*. 1997;21 Suppl 3:S12-23.
  - Ghosal S, Heron N, Mason K, Jordan K. Efficacy of orlistat in type 2 diabetes: A systematic review and meta-analysis. *BJGP Open*. 2025;9(4):BJGPO.2025.0058. <https://doi.org/10.3399/BJGPO.2025.0058>.
  - Filippatos TD, Derdemezis CS, Gazi IF, et al. Orlistat-associated adverse effects and drug interactions. *Drug Saf*. 2008;31(1):53-65. <https://doi.org/10.2165/00002018-200831010-00005>.
  - Caruso I, Cignarelli A, Sorice GP, Perrini S, Giorgino F. Incretin-based therapies for the treatment of obesity-related diseases. *NPJ Metabolic Health and Disease*. 2024;2(1):31. <https://doi.org/10.1038/s44324-024-00030-5>.
  - Zheng Z, Zong Y, Ma Y, et al. Glucagon-like peptide-1 receptor: Mechanisms and advances in therapy. *Signal Transduction and Targeted Therapy*. 2024;9(1):234. <https://doi.org/10.1038/s41392-024-01931-z>.
  - Pi-Sunyer X, Astrup A, Fujioka K, et al. A randomized, controlled trial of 3.0 mg of liraglutide in weight management. *N Engl J Med*. 2015;373(1):11-22. <https://doi.org/10.1056/NEJMoa1411892>.
  - Rubino DM, Greenway FL, Khalid U, et al. Effect of weekly subcutaneous semaglutide vs daily liraglutide on body weight in adults with overweight or obesity without diabetes: The step 8 randomized clinical trial. *JAMA*. 2022;327(2):138-50. <https://doi.org/10.1001/jama.2021.23619>.
  - Jastreboff AM, Kaplan LM, Frías JP, et al. Triple-hormone-receptor agonist retatrutide for obesity - a phase 2 trial. *N Engl J Med*. 2023;389(6):514-26. <https://doi.org/10.1056/NEJMoa2301972>.
  - Halsey G. Retatrutide achieves up to 28.7% weight loss and marked knee pain reduction in phase 3 triumph-4 trial. New York, NY, USA: Haymarket Media, Patient Care Online; 2025. Available from: <https://www.patientcareonline.com/view/retatrutide-achieves-up-to-28-7-weight-loss-and-marked-knee-pain-reduction-in-phase-3-triumph-4-trial>. Accessed 9 January 2026.
  - Van Eenige R, Sánchez-López E, Hoekstra AT, et al. Inhibition of endocannabinoid synthesis enzymes dagl and nape-pld transiently lowers body weight and alters glucose homeostasis during a high-fat diet challenge in mice. *European Journal of Endocrinology*. 2025;193(5):S18-26. <https://doi.org/10.1093/ejendo/lvaf212>.
  - Knop FK, Kunos G, Dicker D, et al. Efficacy and safety of monlunabant in adults with obesity and metabolic syndrome: A double-blind, randomised, placebo-controlled, phase 2a trial. *Lancet Diabetes & Endocrinology*. 2025;13(11):911-23. [https://doi.org/10.1016/S2213-8587\(25\)00216-5](https://doi.org/10.1016/S2213-8587(25)00216-5).
  - Cohen Y, Kolodziej A, Morningstar M. Seventeen years since rimonabant's downfall: Re-assessing its suicidality risk profile. *Obesity*. 2024;32(7):1235-44. <https://doi.org/10.1002/oby.24019>.
  - Kumari P, Dvoráček S, Enos MD, et al. Structural mechanism of cb1r binding to peripheral and biased inverse agonists. *Nature Communications*. 2024;15(1):10694. <https://doi.org/10.1038/s41467-024-54206-0>.